

# Cosmogenic dating and complex patterns of glacier advances during the Lateglacial in the Chagan-Uzun Valley, Russian Altai

Natacha Gribenski<sup>1,2</sup>, Sven Lukas<sup>3</sup>, Krister N. Jansson<sup>1,2</sup>, Arjen P. Stroeven<sup>1,2</sup>, Frank Preusser<sup>4</sup>, Jonathan M. Harbor<sup>1,2,5</sup>, Robin Blomdin<sup>1,2</sup>, Mikhail N. Ivanov<sup>6</sup>, Jakob Heyman<sup>7</sup>, Dmitry Petrakov<sup>6</sup>, Alexei Rudoy<sup>8</sup>, Tom Clifton<sup>9</sup>, Nathaniel A. Lifton<sup>5,9</sup>, Marc W. Caffee<sup>5,9</sup>

<sup>1</sup>Department of Physical Geography, Stockholm University, Sweden. <sup>2</sup>Bolin Centre for Climate Research, Stockholm University, Sweden. <sup>3</sup>School of Geography, Queen Mary University of London, UK.

<sup>4</sup>Institute of Earth and Environmental Sciences - Geology, University of Freiburg, Germany. <sup>5</sup>Department of Earth, Atmospheric, and Planetary Sciences, Purdue University, USA. <sup>6</sup>Cryolithology and Glaciology Department, Moscow State University, Russia. <sup>7</sup>Department of Earth Sciences, University of Gothenburg, Sweden. <sup>8</sup>Department Geology and Geography, National Research Tomsk State University, Russia, <sup>9</sup>Department of Physics and Astronomy, and Purdue Rare Isotope Measurement Laboratory (PRIME Lab), Purdue University, USA

Corresponding author e-mail: [Natacha.gribenski@natgeo.su.se](mailto:Natacha.gribenski@natgeo.su.se)

## Abstract

Over the last decades, numerous paleoglacial reconstructions have been carried out in Central Asian mountain ranges because glaciers in this region are sensitive to climate change, and thus their associated glacial deposits can be used as proxies for paleoclimate inference. However, non-climatic factors can complicate the relationship between glacier fluctuation and climate change. Careful investigations of the geomorphological and sedimentological context are therefore required to understand the mechanisms behind glacier retreat and expansion.

In this study we present the first detailed paleoglacial reconstruction of the Chagan Uzun Valley, located in the Russian Altai. This reconstruction is based on detailed geomorphological mapping, sedimentological logging, and *in situ* cosmogenic <sup>10</sup>Be and <sup>26</sup>Al surface exposure dating of glacially transported boulders.

The Chagan Uzun Valley includes extensive lobate moraine belts (>100 km<sup>2</sup>) deposited in the intramontane Chuja basin, reflecting a series of pronounced former glacial advances. Observation of “hillside-scale” folding and extensive faulting of pre-existing soft sediments within the outer moraine belts, together with the geomorphology, indicate that these moraine belts were formed during glacier-surge like events. In contrast, the inner (up-valley) glacial landforms of the Chagan Uzun valley indicate that they were deposited by temperate alpine glaciers at balance velocity during recessional phases. Cosmogenic ages associated with the outermost, innermost and intermediary stages, all indicate deposition times clustered around 19.2 ka, although the <sup>10</sup>Be ages of the outermost margin are likely slightly underestimated due to brief episode of glacial lake water coverage. Such close deposition timings are consistent with periods of fast or surge advances, followed by active glacier retreat.

This is the first study reporting surge-like behaviour of former glaciers in the Altai mountain range, supported by detailed geomorphological and sedimentological evidences. Such findings are crucial for paleoclimate inference, as the surge-related features cannot be attributed to a glacier system in equilibrium with the contemporary climate, and cannot be interpreted with traditional ELA reconstructions. This study also highlights the complexity of establishing robust paleoglacial chronologies in highly dynamic environments, with interactions between glacial events and the formation and drainage of lakes.